

## Literature Cited

- COLLETTE, B. B.  
1978. II. Adaptations and systematics of the mackerels and tunas. In G. D. Sharp and A. E. Dizon (editors), *The physiological ecology of tunas*, p. 7-39. Acad. Press, N.Y.
- HARADA, T.  
1978. Recent tuna culture research in Japan. 5th International Ocean Development Conference, Keidanren Kaikan, Tokyo, September 25-29, 1978. Preprints (I), Session C-1, p. C1-55-C1-64.
- LEONG, R.  
1977. Maturation and induced spawning of captive Pacific mackerel, *Scomber japonicus*. *Fish. Bull.*, U.S. 75:205-211.
- SHEHADEH, Z. H., C.-M. KUO, AND K. K. MILISEN.  
1973. Validation of an *in vivo* method for monitoring ovarian development in the grey mullet (*Mugil cephalus* L.). *J. Fish Biol.* 5:489-496.
- SŌGŌ.  
1979. Bluefin tuna spawn in captivity—World's first record of artificial fertilization and hatching of bluefin tuna. [In Jpn.] *Sōgō*, June 27, 1979. (Engl. transl. by T. Otsu, 1979, 2 p., Transl. No. 37; available Southwest Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96812.)
- STEVENS, R. E.  
1966. Hormone-induced spawning of striped bass for reservoir stocking. *Prog. Fish-Cult.* 28:19-28.
- THOMAS, A. E.  
1975. Marking channel catfish with silver nitrate. *Prog. Fish-Cult.* 37:250-252.
- UEYANAGI, S.  
1978. Recent tuna culture research in Japan. 5th International Ocean Development Conference, Keidanren Kaikan, Tokyo, September 25-29, 1978. Preprints (I), Session C-1, p. C1-23-C1-30.
- YASUTAKE, H., G. NISHI, AND K. MORI.  
1973. Artificial fertilization and rearing of bigeye tuna (*Thunnus obesus*) on board, with morphological observations on embryonic through to early post-larval stage. [In Jpn., Engl. abstr.] *Bull. Far Seas Fish. Res. Lab. (Shimizu)* 8:71-78.

CALVIN M. KAYA

Southwest Fisheries Center Honolulu Laboratory  
National Marine Fisheries Service, NOAA  
Honolulu, Hawaii  
Present address: Department of Biology  
Montana State University  
Bozeman, MT 59717

ANDREW E. DIZON  
SHARON D. HENDRIX

Southwest Fisheries Center Honolulu Laboratory  
National Marine Fisheries Service, NOAA  
Honolulu, HI 96812

## TROPHIC IMPORTANCE OF SOME MARINE GADIDS IN NORTHERN ALASKA AND THEIR BODY-OTOLITH SIZE RELATIONSHIPS

Natural marine ecosystems are being subjected to ever increasing human-induced stresses, including expanding commercial fisheries and activities associated with the exploration and development of offshore petroleum resources. Numerous studies of the food habits and trophic interactions of marine vertebrate consumers have been undertaken in Alaska during the last 5 yr in response to increased demand for multispecies approaches in fishery management plans and the legal requirement for environmental assessments prior to petroleum development. Through these and other studies the importance of three species—walleye pollock, *Theragra chalcogramma*, saffron cod, *Eleginus gracilis*, and Arctic cod, *Boreogadus saida*—in Arctic and subarctic ecosystems has become increasingly apparent (Klumov 1937; Andriyashev 1954; Lowry and Frost in press; Pereyra et al.<sup>1</sup>). These species are widespread and locally abundant, are major secondary consumers, and are important prey of other species (Table 1).

Walleye pollock are found throughout the North Pacific and in greatest abundance along the continental shelf break of the Bering Sea. Abundance decreases rapidly north of St. Matthew Island, and they are caught only rarely north of Bering Strait (Pereyra et al. footnote 1). The species supports a commercial fishery of almost 1 million t annually, one of the largest in the world. Walleye pollock form a major portion of the diet of all pinnipeds in the southern Bering Sea, except bearded seals and walruses, and are eaten by at least 4 species of cetaceans, 13 species of seabirds, and 10 species of fishes in that area.

Saffron cod occur in the eastern Bering and Chukchi Seas and throughout the western Arctic Ocean (Andriyashev 1954). They are also present, but less abundant, in the Beaufort Sea. Saffron cod are utilized for food by coastal Eskimos. They make up a major portion of the diet of ringed and spotted seals and white whales in the northern Bering and southern Chukchi Seas. They are also

<sup>1</sup>Pereyra, W. T., J. E. Reeves, and R. G. Bakkala. 1976. Demersal fish and shellfish resources of the eastern Bering Sea in the baseline year 1975. Processed rep., 619 p. Northwest and Alaska Fisheries Center, National Marine Fisheries Service, NOAA, 2725 Montlake Boulevard E., Seattle, WA 98112.

TABLE 1.—Marine mammals, birds, and fishes reported to eat walleye pollock, saffron cod, and Arctic cod.

Species	Walleye pollock	Saffron cod	Arctic cod	Species	Walleye pollock	Saffron cod	Arctic cod
<b>Marine mammals:</b>							
Northern fur seal, <i>Callorhinus ursinus</i>	9			Tufted puffin, <i>Lunda cirrhata</i>	21		12
Steller sea lion, <i>Eumetopias jubatus</i>	3, 10, 31			Horned puffin, <i>Fratercula corniculata</i>	21		12
Pacific harbor seal, <i>Phoca vitulina richardsi</i>	23, 24, 31			Kittlitz's murrelet, <i>Brachyramphus brevirostre</i>			15
Spotted seal, <i>P. largha</i>	22, 23, 31	32	22, 23	Parakeet auklet, <i>Cyclorhynchus psittaculus</i>	21		15
Ribbon seal, <i>P. fasciata</i>	14, 23, 31	14, 23	22, 23	Least auklet, <i>Aethia pusilla</i>	21		
Ringed seal, <i>P. hispida</i>	32	11, 23	2, 6, 11, 23	Arctic tern, <i>Sterna paradisaea</i>			4
Bearded seal, <i>Erignathus barbatus</i>	31, 32	8, 23	2, 11, 32	Fulmar, <i>Fulmarus glacialis</i>	21		28
Fin whale, <i>Balaenoptera physalus</i>	5, 13	5	1, 5	Shearwaters, <i>Puffinus</i> spp.			15
Minke whale, <i>B. acutorostrata</i>	5, 25	5	1, 5	Pelagic cormorant, <i>Phalacrocorax pelagicus</i>			12, 15
Sei whale, <i>B. borealis</i>	5		5	Red-faced cormorant, <i>P. urile</i>	21		
Humpback whale, <i>Megaptera novaengliae</i>	5	5	1, 5	Red-throated loon, <i>Gavia stellata</i>			15, 19
White whale, <i>Delphinapterus leucas</i>		5, 25	1, 2, 25	Jaegers, <i>Stercorarius</i> spp.			1
Harp seal, <i>Phoca groenlandica</i>			1	<b>Fishes:</b>			
Narwhal, <i>Monodon monoceros</i>			1, 2, 18	Atlantic cod, <i>Gadus morhua</i>			1
Harbor porpoise, <i>Phocoena phocoena</i>		25		Pacific cod, <i>G. macrocephalus</i>	4, 26, 32		
Polar bear, <i>Ursus maritimus</i>			1	Walleye pollock, <i>Theragra chalcogramma</i>	26, 32		
<b>Birds:</b>				Saffron cod, <i>Eleginus gracilis</i>		4	4
Glaucous gull, <i>Larus hyperboreus</i>	20		12	Pacific halibut, <i>Hippoglossus stenolepis</i>	29		
Herring gull, <i>L. argentatus</i>			12	Greenland halibut, <i>Reinhardtius hippoglossoides</i>	26, 32		
Sabine's gull, <i>Xema sabini</i>			15	Sablefish, <i>Anoploploma fimbria</i>	26		
Ross's gull, <i>Rhodostethia rosea</i>			17	Flathead sole, <i>Hippoglossoides elassodon</i>	29, 32		
Ivory gull, <i>Pagophila eburnea</i>	20		17	American plaice, <i>H. platessoides</i>			1
Black-legged kittiwake, <i>Rissa tridactyla</i>	20, 21	30	1, 12, 30	Arrowtooth flounder, <i>Atheresthes stomias</i>	26, 29		
Red-legged kittiwake, <i>R. brevirostris</i>	20, 21			Snailfish, <i>Liparis</i> sp.	32		
Common murre, <i>Uria aalge</i>	16, 21	30	7, 21, 30	Eelpout, <i>Lycodes</i> spp.	32		
Thick-billed murre, <i>U. lomvia</i>	16, 21	30	7, 28, 30	Sculpins, <i>Icelus spiniger</i> , <i>Myoxocephalus</i> spp.	32	32	
Black guillemot, <i>Cephus grylle</i>	20		1, 12, 28	Sheefish, <i>Stenodus leucichthys</i>		32	
Pigeon guillemot, <i>C. columba</i>			12	Arctic char, <i>Salvelinus alpinus</i>		32	27, 28
				Atlantic salmon, <i>Salmo salar</i>			1
1. Klumov 1937	10. Fiscus and Baines 1966	18. Mansfield et al. 1975					
2. Vibe 1950	11. Johnson et al. 1966	19. Bergman and Derksen 1977					
3. Wilke and Kenyon 1952	12. Swartz 1966	20. Divoky in press					
4. Andriyashev 1954	13. Nemoto 1970	21. Hunt et al. in press					
5. Tomilin 1957	14. Fedoseev and Bukhtiyarov 1972	22. Frost and Lowry 1980					
6. McLaren 1958	15. Watson and Divoky 1972	23. Lowry and Frost in press					
7. Tuck 1960	16. Ogi and Tsujita 1973	24. Pitcher 1980					
8. Kenyon 1962	17. Divoky 1976	25. Frost and Lowry in press					
9. Fiscus et al. 1964							
26. Pereyra et al. (text footnote 1).	report of principal investigators. Vol. I, p. 33-107. Environ. Res. Lab., Boulder, Colo.						
27. Bendock, T. N. 1977. Beaufort Sea estuarine fishery study. In Environmental assessment of the Alaskan continental shelf, annual reports of principal investigators for the year ending March 1977. Vol. VIII, p. 320-365. Environ. Res. Lab., Boulder, Colo.							
28. Bain, H., and A. D. Sekerak. 1978. Aspects of the biology of arctic cod, <i>Boreogadus saida</i> , in the central Canadian arctic. Report for Polar Gas Project by LGL Ltd., Toronto, Ontario, 104 p.							
29. Smith, R. L. 1978. Food and feeding relationships in the benthic and demersal fishes of the Gulf of Alaska and Bering Sea. In Environmental assessment of the Alaskan continental shelf, final							
	30. Springer, A. M., and D. G. Roseneau. 1978. Ecological studies of colonial seabirds at Cape Thompson and Cape Lisburne, Alaska. In Environmental assessment of the Alaskan continental shelf, annual reports of principal investigators for the year ending March 1978. Vol. II, p. 839-960. Environ. Res. Lab., Boulder, Colo.						
	31. Lowry, L. F., K. J. Frost, and J. J. Burns. 1979. Potential resource competition in the southeastern Bering Sea: Fisheries and phocid seals. Proc. 29th Alaska Sci. Conf., p. 287-296.						
	32. Frost and Lowry unpubl. data.						

prey of other cetaceans and numerous birds and fishes.

Arctic cod are circumpolar in Arctic waters extending south to at least lat. 60° N on the Alaska coast, typically in association with sea ice (Andriyashev 1954). They are a species of key trophic importance upon which many other far northern marine consumers depend entirely for a major portion of their yearly nutritional requirements. They are eaten by at least 12 species of marine mammals, 20 species of birds, and 5 species of fishes. Arctic cod are especially important because in the areas and at the times when they are abundant they are the only forage fishes present.

Investigations of food habits of marine animals almost invariably involve analysis of stomach contents. Morrow (1979) published preliminary keys to otoliths of 16 families of fishes found in Alaskan waters including the Gadidae, whereby fishes eaten by predators can be identified from otoliths even after soft parts and bones have been digested. In most instances the size of the fish or meal can also be determined from otoliths through back calculation of fish length and/or weight from various measurements of otolith size (Morrow 1951; Templeman and Squires 1956; Southward 1962; Gjosaeter 1973).

In this paper we present relationships of otolith length to fish length and weight for pollock, saffron cod, and Arctic cod of the Bering, Chukchi, and Beaufort Seas.

#### Methods

Samples of fishes were obtained by otter trawling in the Bering, Chukchi, and Beaufort Seas (Table 2). Soon after capture all fishes were identified, weighed to the nearest 0.1 g, and fork length measured to the nearest millimeter. The sagittal otoliths were removed and length and width measured to the nearest 0.1 mm with vernier calipers.

When otolith lengths and widths were plotted against fish lengths as scatter diagrams, the relationship between otolith length and fish length was found to be less variable than that of otolith width and fish length. For this reason otolith length was taken as the criterion for otolith size and used in subsequent calculations. Casteel (1976) discussed in detail the reasons for using length as the best measure of otolith size.

We chose a double regression method for relating otolith size to fish size (Fitch and Brownell 1968; Casteel 1976). For each species the relationships of otolith length to fish length and fish length to fish weight were calculated. In cases where two equations were required to fit a single relationship, the inflection point was determined by iteration. The specified inflection point was varied by increments of 0.1 and the pair of equations which minimized the combined deviation was selected.

#### Results and Discussion

Regressions of fish fork length on otolith length differed markedly among the three species. Those of walleye pollock and saffron cod formed two distinct straight-line sections each, with inflection points at otolith lengths of 10 mm in walleye pollock (fish length 22 cm) and 8.5 mm in saffron cod (fish length 15 cm) (Figures 1, 2). The regression for Arctic cod was rectilinear over the range of samples (Figure 3).

Several sources of error are possible when estimating the size of a fish from its otoliths, among which are normal variability in the ratio of fish length to otolith length and differences in lengths of left and right otoliths of the same fish. The calculated regression coefficients show that such variability is quite small. Deviation between actual measured and calculated fish lengths was usually <5%. Since food habits studies deal with

TABLE 2.—Sources of Alaskan marine gadids measured to determine otolith length-fish size relationships. T = *Theragra chalcogramma*; E = *Eleginus gracilis*; B = *Boreogadus saida*.

Vessel and cruise no.	Date	Area	Depth range (m)	Trawls (no.)	Species
NOAA <sup>1</sup> Ship <i>Surveyor</i> (RP-4-SU-76A1&II)	Mar.-Apr. 1976	Bering	79-173	39	T
NOAA Ship <i>Discoverer</i> (RP-4-DI-76BIII)	Aug. 1976	Bering/Chukchi	18-55	18	B, E
USCGC <sup>2</sup> <i>Glacier</i> (AWS76)	Aug. 1976	Beaufort	40-123	2	B
NOAA Ship <i>Miller Freeman</i> (RD-4-MF-76BII)	Oct. 1976	Bering	15-55	75	B, E
NOAA Ship <i>Surveyor</i> (RD-4-SU-77AII, III)	Mar.-Apr. 1977	Bering	28-150	45	T, E
NOAA Ship <i>Discoverer</i> (RD-4-DI-77AVI)	May-June 1977	Bering	30-150	36	B, T
NOAA Ship <i>Surveyor</i> (RD-4-SU-77BII)	June-July 1977	Bering/Chukchi	13-57	17	B, E
USCGC <i>Glacier</i> (AWS77II)	Aug.-Sept. 1977	Chukchi/Beaufort	31-400	33	B
ADF&G <sup>3</sup> skiff (Shishmaref 78)	Mar. 1978	Chukchi	5-10	5	E
NOAA Ship <i>Surveyor</i> (RP-4-SU-78AV, VI)	May-June 1978	Bering	17-210	78	T, E

<sup>1</sup>National Oceanic and Atmospheric Administration.

<sup>2</sup>Alaska Department of Fish and Game.

<sup>3</sup>United States Coast Guard Cutter.




FIGURE 1.—Scatter diagram and regression lines and equations of otolith length against fish fork length for *Theragra chalcogramma*.




FIGURE 2.—Scatter diagram and regression lines and equations of otolith length against fish fork length for *Eleginus gracilis*.

mixed collections of otoliths, the cumulative importance of these differences should be minimal.

The relationships between fish lengths and weights of the three species were best fit by exponential equations of the form:  $\text{weight} = a(\text{length})^b$  (Table 3). These relationships may vary somewhat with time of year, geographic location, sex, reproductive status, or fullness of stomach. Variation is probably most pronounced in sexually mature individuals with mature reproductive products, a condition which persists for only a few months of the year. Since small (juvenile) fishes are eaten by most marine mammals (Frost and Lowry 1980), birds (Hunt et al. in press), and other fishes

(Frost and Lowry unpubl. data), this is probably a small source of error. Significant differences in weight-at-length by sex and geographic area were found for Arctic and saffron cods by Wolotira et al.<sup>2</sup> but they justified use of a single regression equation since the differences were small (3-7%). Similar differences have been noted for walleye pollock (Bakkala and Smith<sup>3</sup>).

Otoliths are valuable indicators of the diet of piscivorous marine consumers. Published keys such as Morrow (1979) allow determination of the species and numbers of fishes represented by otoliths in stomachs, intestines, or scats. By using the relationships between otolith size and body

TABLE 3.—Length-weight relationships observed for walleye pollock, saffron cod, and Arctic cod in the Bering, Chukchi, and Beaufort Seas ( $\text{weight} = a(\text{length})^b$ ).

Species	Number sampled	Range in fork length (cm)	a	b	Regression coefficient (r)
Walleye pollock	109	6-57	0.0077	2.906	0.998
Saffron cod	104	6-29	.0050	3.095	.991
Arctic cod	118	7-21	.0018	3.500	.987

<sup>2</sup>Wolotira, R. J., Jr. 1977. Demersal fish and shellfish resources of Norton Sound, the southeastern Chukchi Sea and adjacent waters in the baseline year 1976. Processed rep., 292 p. Northwest and Alaska Fisheries Center, National Marine Fisheries Service, NOAA, 2725 Montlake Boulevard E., Seattle, WA 98112.

<sup>3</sup>Bakkala, R. G., and G. B. Smith. 1978. Demersal fish resources of the eastern Bering Sea: Spring 1976. Processed rep., 233 p. Northwest and Alaska Fisheries Center, National Marine Fisheries Service, NOAA, 2725 Montlake Boulevard E., Seattle, WA 98112.




FIGURE 3.—Scatter diagram and regression lines and equations of otolith length against fish fork length for *Boreogadus saida*.

size it is possible to obtain additional information such as sizes and quantities of fishes eaten by consumers (Frost and Lowry 1980).

#### Acknowledgments

Many people assisted us in the collection of samples, especially Larry M. Shults who spent many long hours sorting through trawls and measuring fish with us, and the officers and crew of the NOAA Ship *Surveyor* who gave unstintingly of their time and energy to make our project a success. Lawrence R. Miller provided invaluable assistance in the computer analysis of our data. We thank J. E. Morrow and anonymous reviewers for their careful review of the manuscript. We are especially indebted to John Fitch for his many helpful suggestions and the moral support he lent throughout preparation of this manuscript. Financial support was provided by the U.S. Bureau of Land Management Outer Continental Shelf Environmental Assessment Program and Federal Aid in Wildlife Restoration Project W-17-9.

#### Literature Cited

- ANDRIYASHEV, A. P.  
1954. Ryby severnykh morei SSSR (Fishes of the northern seas of the USSR). Izd. Akad. Nauk SSSR, Keys to the fauna of the USSR 53. (Translated from Russ. by Isr. Program Sci. Transl., 1964, 617 p.; available Natl. Tech. Inf. Serv., Springfield, Va., as OTS 63-11160.)
- BERGMAN, R. D., AND D. V. DERKSEN.  
1977. Observations on arctic and red-throated loons at Storkersen Point, Alaska. *Arctic* 30:41-51.
- CASTEEL, R. W.  
1976. Fish remains in archaeology and paleo-environmental studies. *Acad. Press, Lond.*, 180 p.
- DIVOKY, G. J.  
1976. The pelagic feeding habits of Ivory and Ross' Gulls. *Condor* 78:85-90.  
In press. Birds and ice relations. In D. W. Hood (editor), *The Eastern Bering Sea Shelf: Oceanography and resources*, Sect. 13.
- FEDOSEEV, G. A., AND YU. A. BUKHTIYAROV.  
1972. Pitaniye tulenei okhotskogo more (The diet of seals of the Okhotsk Sea). [Abstr.] In V. A. Arsen'ev, V. Bel'kovich, V. A. Zemskii, B. A. Zenkovich, V. E. Sokolov, and K. K. Chapskii (editors), *Theses of Works Fifth All-Union Conference on Studies of Marine Mammals, Makhachkala, USSR, Part 1*, p. 110-112.
- FISCUS, C. H., AND G. A. BAINES.  
1966. Food and feeding behavior of Steller and California sea lions. *J. Mammal.* 47:195-200.
- FISCUS, C. H., G. A. BAINES, AND F. WILKE.  
1964. Pelagic fur seal investigations Alaska Waters, 1962. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 475, 59 p.
- FITCH, J. E., AND R. L. BROWNELL, JR.  
1968. Fish otoliths in cetacean stomachs and their importance in interpreting feeding habits. *J. Fish. Res. Board Can.* 25:2561-2574.
- FROST, K. J., AND L. F. LOWRY.  
1980. Feeding of ribbon seals (*Phoca fasciata*) in the Bering Sea in spring. *Can. J. Zool.* 58:1601-1607.  
In press. Foods and trophic relationships of cetaceans in the Bering Sea. In D. W. Hood (editor), *The Eastern Bering Sea Shelf: Oceanography and resources*, Sect. 10.
- GJOSAETER, J.  
1973. Preliminary results of Norwegian polar cod investigations 1970-1972. *Int. Counc. Explor. Sea Rep.*, 23 p.
- HUNT, G. L., JR., B. BURGESSON, AND G. SANGER.  
In press. Feeding ecology of seabirds of the eastern Bering Sea. In D. W. Hood (editor), *The Eastern Bering Sea Shelf: Oceanography and resources*, Sect. 11.
- JOHNSON, M. L., C. H. FISCUS, B. T. OSTENSON, AND M. L. BARBOUR.  
1966. Marine mammals. In N. J. Wilimovsky and J. N. Wolfe (editors), *Environment of the Cape Thompson region, Alaska*, p. 897-924. U.S. At. Energy Comm., Wash., D.C.
- KENYON, K. W.  
1962. Notes on the phocid seals at Little Diomed Island, Alaska. *J. Wildl. Manage.* 26:380-387.
- KLUMOV, S. K.  
1937. Saika (*Boreogadus saida* (Lepech.)) i ee znachenie dlja nekotorykh zhiznennik protzesov arktiki (Morue polaire (*Boreogadus saida*) et son importance pour cer-

- tains procès vitaux de l'Arctique.) [In Russ., Fr. abstr.] *Izv. Akad. Nauk SSSR* No. 1, p. 175-181.
- LOWRY, L. F., AND K. J. FROST.  
In press. Feeding and trophic relationships of phocid seals and walruses in the eastern Bering Sea. In D. W. Hood (editor), *The Eastern Bering Sea Shelf: Oceanography and resources*, Sect. 10.
- MANSFIELD, A. W., T. G. SMITH, AND B. BECK.  
1975. The narwhal, *Monodon monoceros*, in eastern Canadian waters. *J. Fish. Res. Board Can.* 32:1041-1046.
- MCLAREN, I. A.  
1958. The biology of the ringed seal (*Phoca hispida* Schreber) in the eastern Canadian arctic. *Fish. Res. Board Can.*, Bull. 118, 97 p.
- MORROW, J. E., JR.  
1951. Studies on the marine resources of southern New England. VIII. The biology of the longhorn sculpin, *Myoxocephalus octodecimspinosus* Mitchell, with a discussion of the southern New England "trash" fishery. *Bull. Bingham Oceanogr. Collect.*, Yale Univ. 13(2), 89 p.  
1979. Preliminary keys to otoliths of some adult fishes of the Gulf of Alaska, Bering Sea, and Beaufort Sea. U.S. Dep. Commer., NOAA Tech. Rep. NMFS CIRC-420, 32 p.
- NEMOTO, T.  
1970. Feeding pattern of baleen whales in the ocean. In J. H. Steele (editor), *Marine food chains*, p. 241-252. Univ. Calif. Press, Berkeley.
- OGI, H., AND T. TSUJITA.  
1973. Preliminary examination of stomach contents of murrets (*Uria* spp.) from the eastern Bering Sea and Bristol Bay, June-August, 1970 and 1971. *Jpn. J. Ecol.* 23:201-209.
- PITCHER, K. W.  
1980. Food of the harbor seal, *Phoca vitulina richardsi*, in the Gulf of Alaska. *Fish. Bull.*, U.S. 78:544-549.
- SOUTHWARD, G. M.  
1962. Photographing halibut otoliths for measuring growth zones. *J. Fish. Res. Board Can.* 19:335-338.
- SWARTZ, L. G.  
1966. Sea-cliff birds. In N. J. Wilimovsky and J. N. Wolfe (editors), *Environment of the Cape Thompson region, Alaska*, p. 611-678. U.S. At. Energy Comm., Wash., D.C.
- TEMPLEMAN, W., AND H. J. SQUIRES.  
1956. Relationship of otolith lengths and weights in the haddock, *Melanogrammus aeglefinus* (L.) to the rate of growth of the fish. *J. Fish. Res. Board Can.* 13:467-487.
- TOMILIN, A. G.  
1957. Kitoobraznye (Cetacea). In V. G. Heptner (editor), *Zveri vostochnoi Evropy i severnoi Azii* (Mammals of the USSR and adjacent countries), Vol. IX. (Translated by Isr. Program Sci. Transl., 1967, 717 p.; available Natl. Tech. Inf. Serv., Springfield, Va., as TT-65-50086.)
- TUCK, L. M.  
1960. The murrets; their distribution, populations and biology; a study of the genus *Uria*. *Can. Wildl. Serv. Rep.*, Ser. 1, 260 p.
- VIBE, C.  
1950. The marine mammals and the marine fauna in the Thule district (northwest Greenland) with observations on ice conditions in 1939-41. *Medd. Grønland* 150(6), 115 p.
- WATSON, G. E., AND G. J. DIVOKY.  
1972. Pelagic bird and mammal observations in the eastern Chukchi Sea, early fall 1970. U.S. Coast Guard Oceanogr. Rep. 50:111-172.
- WILKE, F., AND K. W. KENYON.  
1952. Notes on the food of the fur seal, sea-lion and harbor porpoise. *J. Wildl. Manage.* 16:396-397.

KATHRYN J. FROST  
LLOYD F. LOWRY

*Alaska Department of Fish and Game*  
1300 College Road  
Fairbanks, AK 99701

CAROLINIAN RECORDS FOR AMERICAN  
LOBSTER, *HOMARUS AMERICANUS*, AND  
TROPICAL SWIMMING CRAB,  
*CALLINECTES BOCOURTI*. POSTULATED  
MEANS OF DISPERSAL

Recent reports of distributional extension for decapod crustaceans occurring along the east coast of the United States include two poorly substantiated records of American lobster, *Homarus americanus* H. Milne Edwards, and none of the tropical swimming crab, *Callinectes bocourti* A. Milne Edwards, from the Carolinas south of Cape Hatteras, N.C. (Williams 1965, 1974 [Carolinas]; Cerase-Vivas and Gray 1966 [Cape Hatteras]; Williams et al. 1968 [North Carolina]; Musick and McEachren 1972 [North Carolina-Virginia]; Milstein et al. 1977 [New Jersey]; Bowen et al. 1979 [Middle Atlantic area]; Herbst, Weston, and Lorman 1979 [Cape Hatteras]; Herbst, Williams, and Boothe 1979 [Capes Hatteras and Lookout]; Wenner and Boesch 1979 [Norfolk Canyon area]; Perschbacher and Schwartz 1979 [North Carolina]). Occurrences of both species in the Carolinas south of Cape Hatteras are documented here along with discussion of their postulated means of dispersal.

Specimens are deposited in the U.S. National Museum of Natural History (USNM), or are living in aquaria at the North Carolina Marine Resources Center, Bogue Banks (NCMRC), and the Hampton Mariners Museum, Beaufort (HMM).

Occurrence of Species

*Homarus americanus*.—Distribution of the American lobster has been given as, "East coast of America from the Strait of Belle Isle, Newfoundland (Canada) to Cape Hatteras, North Carolina (U.S.A.)," at depths of 0-480 m, usually 4-50 m (Holthuis 1974). Reported occurrences of this species south of Cape Hatteras are: one caught in a